

Age-standardized (world) incidence (per 100,000) and cumulative (0-74) incidence (percent) rates and standard errors

Nose, sinuses etc. (C30-31) (contd)

	Male					Female				
	Cases	ASR(W)	CUM 0-74			Cases	ASR(W)	CUM 0-74		
Oceania (contd)										
*Australia, Northern Territory : Non-Indigenous	1	0.2	0.15	0.02	0.02	0	—	0.00	—	0.00
*Australia, Northern Territory : Indigenous	2	1.4	1.06	0.16	0.12	0	—	0.00	—	0.00
*Australia, Queensland	109	0.6	0.06	0.07	0.01	77	0.4	0.05	0.04	0.01
Australia, South Australia	47	0.6	0.10	0.07	0.01	26	0.3	0.07	0.03	0.01
Australia, Tasmania	25	1.0	0.22	0.11	0.03	9	0.4	0.14	0.04	0.02
*Australia, Victoria	152	0.6	0.05	0.06	0.01	80	0.3	0.04	0.04	0.01
*Australia, Western Australia	64	0.7	0.09	0.07	0.01	39	0.4	0.07	0.05	0.01
*New Caledonia	6	0.8	0.33	0.10	0.04	1	0.1	0.07	0.00	0.00
New Zealand	92	0.5	0.06	0.06	0.01	50	0.3	0.04	0.03	0.00
*New Zealand : Maori	12	0.7	0.21	0.10	0.03	4	0.2	0.11	0.03	0.01
*New Zealand : Pacific peoples	7	1.1	0.42	0.16	0.07	1	0.1	0.13	0.01	0.01
*New Zealand : Other	69	0.5	0.06	0.05	0.01	40	0.2	0.04	0.03	0.01
USA, Hawaii	44	0.7	0.12	0.10	0.02	17	0.2	0.06	0.03	0.01
USA, Hawaii : White	15	0.7	0.20	0.12	0.03	5	0.2	0.11	0.04	0.02
*USA, Hawaii : Japanese	7	0.4	0.16	0.04	0.02	4	0.0	0.02	0.00	0.00
*USA, Hawaii : Chinese	5	2.8	1.38	0.28	0.12	1	0.4	0.42	0.03	0.03
*USA, Hawaii : Hawaiian	5	0.6	0.25	0.08	0.04	2	0.2	0.15	0.02	0.02
*USA, Hawaii : Filipino	8	1.0	0.36	0.12	0.05	2	0.2	0.14	0.03	0.02

*Important. See note on population page.